

Drug Facts

Daytime QuickMelts

Active Ingredient (per tablet)	Purpose
Zincum Gluconicum 1X	Cold Remedy

Uses

- to reduce the duration of the common cold
- reduces the severity of cold symptoms: cough, sore throat, nasal congestion, post nasal drip and/or hoarseness

Warnings

Ask a doctor before use if you

- are taking minocycline, doxycycline, tetracycline or are on coumadin therapy, zinc treatment may inhibit the absorption of these medicines

Stop use and ask your healthcare practitioner if symptoms persist beyond 7 days.

Cold-EEZE® QuickMelts® Daytime is formulated to reduce the duration of the common cold and its symptoms and is insufficient treatment for Influenza or Allergies.

If pregnant or breastfeeding, ask a healthcare practitioner before use. **Keep out of reach of children.**

Directions

- Take Cold-EEZE® Daytime QuickMelts® when you have cold symptoms
- Adults and Children 12 years and over:
 - Dissolve entire tablet in mouth. Do not chew. Do not swallow whole.
 - Take 1 tablet at the onset of symptoms.
 - Repeat every 2-4 hours as needed until all symptoms subside.
 - Do not eat or drink for 15 minutes after use, otherwise, drink plenty of fluids.
 - Recommended daily dose is 6 tablets for adults and 4 tablets for ages 12-17 years.
 - For children under 12 years of age, consult a healthcare practitioner before use.

Other information

- Do not take Cold-EEZE® QuickMelts® on an empty stomach to avoid possible minor stomach upset.
- Store in a cool and dry place.

Inactive Ingredients

crospovidone, FD&C Red #40, glycine, magnesium stearate, malic acid, mannitol, microcrystalline cellulose, natural mixed berry flavor, natural menthol flavor, polyethylene glycol, silicon dioxide, sodium lauryl sulfate, sodium starch glycolate, stevia, sucralose, talc.

Drug Facts

Nighttime QuickMelts

Active ingredient (per tablet)	Purpose
Zincum Gluconicum 1X	Cold Remedy
Valeriana Officinalis 1X	Sleep Disorder Remedy
Chamomile (Matricaria recutita) 2X	Relaxation and Calming

Uses

Take 1 Nighttime QuickMelt 1 hour before bedtime per Directions

- to reduce the duration of the common cold.
- reduces the severity of cold symptoms: cough, sore throat, nasal congestion post nasal drip and/or hoarseness.
- helps to relax and sleep easier to get rested.

Warnings

Ask a doctor before use if you

- are taking fluvoxamine (Luvox), minocycline, doxycycline, tetracycline or are on Coumadin therapy, as zinc treatment may inhibit the absorption of these medicines.
- are under medical supervision and/or are taking tranquilizers, sedatives or anti-seizure medicines.
- do not take concomitantly with loperamide.

Stop use and ask your healthcare practitioner if symptoms persist beyond 7 days.

Cold-EEZE® Nighttime QuickMelts® is a combination product formulated to help you to sleep easier and to reduce the duration of the common cold and its symptoms and is insufficient treatment for Influenza or Allergies.

Uses

Take 1 Nighttime QuickMelt 1 hour before bedtime per Directions

- to reduce the duration of the common cold.
- reduces the severity of cold symptoms: cough, sore throat, nasal congestion post nasal drip and/or hoarseness.
- helps to relax and sleep easier to get rested.

Warnings

Ask a doctor before use if you

- are taking fluvoxamine (Luvox), minocycline, doxycycline, tetracycline or are on Coumadin therapy, as zinc treatment may inhibit the absorption of these medicines.
- are under medical supervision and/or are taking tranquilizers, sedatives or anti-seizure medicines.
- do not take concomitantly with loperamide.

Stop use and ask your healthcare practitioner if symptoms persist beyond 7 days.

Cold-EEZE® Nighttime QuickMelts® is a combination product formulated to help you to sleep easier and to reduce the duration of the common cold and its symptoms and is insufficient treatment for Influenza or Allergies.

Do not take Cold-EEZE® Nighttime QuickMelts® while operating a motor vehicle or machinery or consuming alcohol.

Allow 6-8 hours for sleep after taking tablet.

If pregnant or breastfeeding, ask a healthcare practitioner before use.

Keep out of reach of children.

Directions

- Take Cold-EEZE® Nighttime QuickMelts® at night when you have cold symptoms.
- Adults:
 - Dissolve entire tablet in mouth. Do not chew. Do not swallow whole.
 - Take 1 Nighttime tablet 1 hour before bedtime.
 - Do not eat or drink for 15 minutes after use, otherwise, drink plenty of fluids.
- Recommended dose is 1 tablet for adults.
- For individuals under the age of 18, consult a healthcare practitioner before use.

Other Information

- Do not take Cold-EEZE® QuickMelts® on an empty stomach to avoid possible minor stomach upset.
- Store in a cool and dry place.

Inactive Ingredients

crospovidone, FD&C Blue #1 & #2, glycine, magnesium stearate, malic acid, maltodextrin, mannitol, microcrystalline cellulose, natural mixed berry flavor, natural menthol flavor, polyethylene glycol, sodium lauryl sulfate, silicon dioxide, sodium starch glycolate,